



(infliximab-dyyb)

INFLECTRA Infusion Orders

Patient Name _____ DOB _____ Gender: M F

Phone _____

DIAGNOSIS: *Please provide IDC-10 Code*

- | | |
|--|---|
| <input type="checkbox"/> _____ Rheumatoid Arthritis (RA) | <input type="checkbox"/> _____ Crohn's Disease |
| <input type="checkbox"/> _____ Psoriatic Arthritis | <input type="checkbox"/> _____ Ulcerative Colitis |
| <input type="checkbox"/> _____ Plaque Psoriasis | <input type="checkbox"/> _____ (other) |
| <input type="checkbox"/> _____ Ankylosing Spondylitis | |

PRE-MEDICATION:

- | | |
|--|---|
| <input type="checkbox"/> Tylenol 1000mg PO | <input type="checkbox"/> Solu-Medrol 125mg IVP |
| <input type="checkbox"/> Cetirizine 10mg PO | <input type="checkbox"/> Solu-Cortef 100mg IVP |
| <input type="checkbox"/> Diphenhydramine 25mg PO | <input type="checkbox"/> Diphenhydramine 25mg IVP |
| <input type="checkbox"/> _____ Other | <input type="checkbox"/> _____ Other |

INFLECTRA ORDERS:

DOSAGE:

- _____ mg/kg (weight based)
- _____ mg (flat dose)
- ROUND TO THE NEAREST 100 mg VIAL**
- Emergency Orders and Treatment per Novello Infusion Anaphylaxis Policy**

PATIENT WEIGHT

(Current, but weigh each visit)

_____ LBS

_____ KG

FREQUENCY:

- At 0,2,6 and every 8 weeks
- At 0,2,6 and every _____ weeks
- Other _____

NOTES:

Nursing Orders

- Assess patient prior to each infusion
- Provide patient/caregiver education related to disease process, therapy, infection control, drug use, side effects and precautions, emergency plan
- Weigh patient every visit and record
- Monitor vital signs initially and throughout infusion per manufacturer recommendations
- Establish and maintain IV access / may use central venous access if appropriate and maintain per Novello policy
- Flush IV with 3ml-5ml of saline before and after every IV medication and PRN
- Administer infusion medication/injection as ordered
- If mild itching, slow down the infusion and monitor patient closely
- If anaphylactic reaction, stop infusion, implement emergency medications, and plan and call 911
- When infusion complete, flush IV with 3ml to 5ml saline flush or 50ml bag of saline and discontinue IV per INS Standards
- Evaluate effectiveness of infused therapy on disease process
- Lab work per physician when ordered

ORDERING PROVIDER:

Signature _____ Printed Name: _____ Date _____

Practice _____ Phone _____ Fax _____

**** PLEASE FAX COMPLETED ORDER TO NOVELLO INFUSION 231-600-7058****