



(ocrelizumab)

# OCREVUS Infusion Orders

Patient Name \_\_\_\_\_ DOB \_\_\_\_\_ Gender:  M  F

Phone \_\_\_\_\_

**DIAGNOSIS:** *Please provide IDC-10 Code*

- \_\_\_\_\_ Multiple Sclerosis
- \_\_\_\_\_ (Other)

**PRE-MEDICATION:**

- Cetirizine 10mg PO
- Tylenol 1000mg PO
- \_\_\_\_\_ Other
- \_\_\_\_\_ Other

**OCREVUS ORDERS:**

**DOSAGE/FREQUENCY:**

- 300mg IV initial does, followed 2 weeks later by a second 300mg IV dose
- Subsequent to first 2 doses, 600mg IV dose every 6 months.
- Emergency Orders and Treatment per Novello Infusion Anaphylaxis Policy**

**PATIENT WEIGHT**

\_\_\_\_\_ LBS  
\_\_\_\_\_ KG

**PRE-MEDICATION PER PRESCRIBING INFORMATION:**

- Solu-medrol 100mg IV 30 minutes prior to each treatment
- Diphenhydramine 50mg PO 30 minutes prior to each treatment

**NOTES:**

**Nursing Orders**

- Assess patient prior to each infusion
- Provide patient/caregiver education related to disease process, therapy, infection control, drug use, side effects and precautions, emergency plan
- Weigh patient every visit and record
- Monitor vital signs initially and throughout infusion per manufacturer recommendations
- Establish and maintain IV access / may use central venous access if appropriate and maintain per Novello policy
- Flush IV with 3ml-5ml of saline before and after every IV medication and PRN
- Administer infusion medication/injection as ordered
- If mild itching, slow down the infusion and monitor patient closely
- If anaphylactic reaction, stop infusion, implement emergency medications, and plan and call 911
- When infusion complete, flush IV with 3ml to 5ml saline flush or 50ml bag of saline and discontinue IV per INS Standards
- Evaluate effectiveness of infused therapy on disease process
- Lab work per physician when ordered

**ORDERING PROVIDER:**

Signature \_\_\_\_\_ Printed Name: \_\_\_\_\_ Date \_\_\_\_\_

Practice \_\_\_\_\_ Phone \_\_\_\_\_ Fax \_\_\_\_\_

\*\*\*\*PLEASE FAX COMPLETED ORDER TO NOVELLO INFUSION 231-600-7058\*\*\*\*